



Short communication

Population structure of *Blackberry yellow vein associated virus*, an emerging crinivirus[☆]Bindu Poudel^a, Sead Sabanadzovic^b, Jozef Bujarski^{c,d}, Ioannis E. Tzanetakis^{a,*}^a Department of Plant Pathology, Division of Agriculture, University of Arkansas, Fayetteville, AR 72701, USA^b Department of Biochemistry, Molecular Biology, Entomology, and Plant Pathology, Mississippi State University, Mississippi State, MS 39762, USA^c Department of Biological Sciences, Northern Illinois University, DeKalb, IL 60115, USA^d Institute of Bioorganic Chemistry, Polish Academy of Sciences, Poznan, Poland

ARTICLE INFO

Article history:

Received 23 May 2012

Received in revised form 9 July 2012

Accepted 12 July 2012

Available online 20 July 2012

Keywords:

Closteroviridae

Recombination

Selection pressure

Evolution

ABSTRACT

Blackberry yellow vein disease (BYVD), a disorder caused by virus complexes, has become a major threat to fresh market blackberry production in the United States. *Blackberry yellow vein associated virus* (BYVaV) is the most prevalent virus in the BYVD complexes; detected in about 50% of samples exhibiting typical disease symptoms. Thirty-four virus isolates infecting wild and cultivated blackberries were collected from several areas with high BYVD incidence. Sequence variability and virus evolution predictions were calculated for four genomic regions coding for six proteins and accounting for about 30% of the virus genome. Nucleotide diversity ranged between 7 and 12%, and all proteins studied were under negative selection. Several isolates were identified as potential recombinants suggesting that recombination might be a driving force behind BYVaV evolution.

© 2012 Elsevier B.V. All rights reserved.

Recent increase in blackberry acreage and expansion to new areas has led to an increased incidence of viruses and the emergence of a new disorder designated as blackberry yellow vein disease (BYVD; Martin et al., 2012). Martin et al. (2004) discovered a new crinivirus named *Blackberry yellow vein associated virus* (BYVaV) in all symptomatic BYVD plants. Further study determined that BYVaV is not the causal agent of the disorder and that BYVD is rather caused by diverse virus complexes (Susaimuthu et al., 2008). Notwithstanding, BYVaV is the most prevalent virus in the complexes; detected in several commercial nurseries scattered around the United States, all fresh market blackberry growing areas, and several wild blackberry populations (Susaimuthu et al., 2007; Poudel, 2011).

BYVaV is a member of the genus *Crinivirus*, family *Closteroviridae* and has a bipartite genome of 15.7 kb in size. The 7.8 kb RNA1 codes for a papain-like protease, methyltransferase, helicase and polymerase, all involved in virus replication; whereas the 7.9 kb RNA2 codes for proteins involved in encapsidation, transmission and movement (Tzanetakis et al., 2006). An isolate from an 'Apache' blackberry from South Carolina has been sequenced (Tzanetakis et al., 2006) and represents the virus type isolate; however, knowledge on additional BYVaV isolates is limited. Although

asymptomatic in single infections (Susaimuthu et al., 2007), BYVaV interactions with other viruses indicate that disease symptoms are elicited upon infection with different viruses (Susaimuthu et al., 2008). It is therefore important to study the virus population structure and set the foundation to better understand BYVaV isolates/strains interactions with other viruses as a part of BYVD development. This communication provides a comprehensive analysis of the population structure of BYVaV with information on 34 isolates collected from vast geographical area that spans several hundred thousand km² (Table 1).

Young, fully developed leaves showing BYVD symptoms were collected from the wild in Arkansas and commercial fields in Arkansas, Georgia, Mississippi, North Carolina, and South Carolina. The Arkansas isolates were studied in more detail as the University of Arkansas has a leading breeding program for fresh market blackberries and infected material from the program could be the virus source of material grown around the world. Samples were processed within 24 h of collection. Total nucleic acid preparations were extracted as described by Tzanetakis et al. (2007) and subjected to reverse-transcription polymerase chain reaction (RT-PCR) with two sets of detection primers (Susaimuthu et al., 2006, 2008), so as to identify the highest possible number of BYVaV isolates. Positive samples were subjected to dsRNA extractions as described in Laney et al. (2011). All regions were amplified from reverse transcribed dsRNA templates essentially as described in Tzanetakis et al. (2005). The designation of the different regions studied, along with nucleotide coordinates are: 1 [polyprotein (1a): RNA1 nt 2384–3583], 2 [RNA-dependent RNA polymerase (RdRp);

[☆] Sequences reported in this communication have been deposited in Genbank and have been assigned the accession numbers JQ56161–JQ56296.

* Corresponding author. Tel.: +1 479 575 3180; fax: +1 479 575 7601.

E-mail address: itzaneta@uark.edu (I.E. Tzanetakis).

Table 1
Isolates used in population structure study of *Blackberry yellow vein associated virus*.

Isolate name	Wild/cultivated	Collection state
AR1	Cultivated	Arkansas
AR2	Cultivated	Arkansas
AR3	Cultivated	Arkansas
AR4	Cultivated	Arkansas
AR5	Cultivated	Arkansas
AR6	Cultivated	Arkansas
AR7	Cultivated	Arkansas
AR8	Wild	Arkansas
AR9	Wild	Arkansas
AR10	Wild	Arkansas
AR11	Wild	Arkansas
AR12	Wild	Arkansas
AR13	Wild	Arkansas
GA1	Cultivated	Georgia
GA2	Cultivated	Georgia
GA3	Cultivated	Georgia
MS1	Cultivated	Mississippi
MS2	Cultivated	Mississippi
MS3	Cultivated	Mississippi
NC1	Cultivated	North Carolina
NC2	Cultivated	North Carolina
NC3	Cultivated	North Carolina
NC4	Cultivated	North Carolina
NC5	Cultivated	North Carolina
NC6	Cultivated	North Carolina
NC7	Cultivated	North Carolina
SC1	Cultivated	South Carolina
SC2	Cultivated	South Carolina
SC3	Cultivated	South Carolina
SC4	Cultivated	South Carolina
SC5	Cultivated	South Carolina
SC6	Cultivated	South Carolina
SC7	Cultivated	South Carolina
SC8	Cultivated	South Carolina
SCO	Type isolate	South Carolina

RNA1 nt 6319–7518], 3 [p4 and part of heat shock protein 70 homolog (HSP70h); RNA2 nt 309–1103] and 4 [part of the coat protein (CP) and minor coat protein (CPm); RNA2 nt 4709–6024] (Fig. 1).

Purified PCR products were cloned onto the TOPO-TA pCR2.1 vector (Invitrogen) and transformed into α -select chemically competent cells (Bioline). Selected recombinant plasmids were sequenced by Functional Biosciences, Inc. (Madison, WI) using universal M13 forward and reverse primers. A minimum of three independent clones were sequenced in both directions to obtain the consensus sequence for each region studied.

Sequence contigs were obtained with CAP3 (Huang and Madan, 1999). Diversity among isolates was studied using Bioedit (Hall, 1999). Selection pressure was calculated using the Synonymous/Non-Synonymous Analysis Program (SNAP; Korber, 2000). The program calculates synonymous/non-synonymous rates based on Nei–Gojobori method (Nei and Gojobori, 1986). The results were further validated by using the codon based Z-test, also implementing the Nei–Gojobori method (Nei and Gojobori, 1986).




Fig. 1. Schematic representation of the BYVaV genomic regions used for the diversity study indicated by left-right arrows. Abbreviations: Pro: papain-like protease; MT: methyltransferase; HEL: helicase; Pol: RNA-dependent RNA polymerase; HSP70h: heat shock protein homolog; CPh: coat protein homolog; CP: coat protein; CPm: minor coat protein.

on MEGA v.5 (Tamura et al., 2011). Phylogenetic and molecular evolutionary analyses also were conducted with MEGA. The maximum likelihood method was used to construct the trees using the nucleotide sequences for the four regions studied implementing the Kimura 2-parameter model (Kimura, 1980). Trees were also constructed for six coding regions using the Poisson model for amino acid substitution (Zuckerland and Pauling, 1965). In both analyses, grouping robustness was determined after 1000 pseudoreplicates. Branches supported by <50% were collapsed as they were considered untrustworthy (Felsenstein, 1985). Recombination analysis was performed on RDP4 which implements eight algorithms to identify recombination sites (Martin et al., 2010). Recombination patterns were considered statistically significant events when detected by four or more programs (Martin et al., 2010).

Four regions, two from each genomic RNA, were analyzed, accounting for about 30% of the BYVaV genome (4512 nt/isolate). It is generally accepted that replication-associated proteins as those coded in RNA1 show a slower evolutionary rate compared to those coded by RNA2, involved in movement and genome protection (Dolja et al., 2006). On the other hand isolates of the monopartite closteroviruses and ampeloviruses tend to show higher nucleotide diversity toward the 5' end of the genome (Roy et al., 2005; Herrera-Isidró et al., 2009; Aguilar et al., 2003) but this is grossly understudied in criniviruses, a gap in knowledge this study aims to address.

The positioning of region 1 (polyprotein) was based on the presence of additions/deletions in the orthologous region of *Beet pseudo-yellows virus* (BPYV), the closest relative to BYVaV identified to date (Tzanetakis and Martin, 2004). Region 2 codes for the RdRp and as in the case of the polyprotein, is less studied in crinivirus population studies. Additional rationale for choosing RdRp is that our data on *Strawberry pallidosis associated virus* indicate significant diversity in this genomic region (Tzanetakis et al., unpublished). BYVaV code for two small hydrophobic peptides with identifiable transmembrane domains near the 5' terminus of RNA2. This has not been observed for any other crinivirus, thus region 3 was designed to evaluate the presence of the peptides and the transmembrane domains in all isolates. The C-termini of the crinivirus coat proteins contain the conserved Arg and Asp residues, found in all filamentous RNA viruses, and are more conserved compared to the N-termini of the proteins (Tzanetakis et al., 2006). For this reason, region 4 was designed to include the CP/CPm junction and determine whether the two areas are under different selection pressures. Several primer set combinations were evaluated to determine the most efficient PCR amplification of the studied regions. Based on those experiments the regions analyzed are slightly altered from the original design. The data in this communication represents the minimum sequence length as amplified from all isolates.

The ratio of non-synonymous substitutions per non-synonymous site (d_{NS}) over synonymous substitutions per synonymous site (d_S) was used as the indicator of protein selection pressure. Coding areas are under positive or diversifying selection when $d_{NS}/d_S > 1$, neutral when $d_{NS}/d_S = 1$, and negative (purifying) when $d_{NS}/d_S < 1$. The d_{NS}/d_S ratio was calculated pairwise for isolates and averaged for a particular region. Region 1, showed stability with maximum nucleotide and amino acid diversity of 8% and 6%, respectively, and the $d_{NS}/d_S = 0.042$. Apparently, region 2 was under stringent purification selection. Although the nucleotide diversity among studied isolates reached 12%, the maximum amino acid diversity was only 5% and $d_{NS}/d_S = 0.021$ (Table 2).

Region 3 covered two coding regions, p4 and the N-terminus of the HSP70h. The maximum nucleotide and amino acid diversity of p4 reached 12% and 11%, respectively. Accordingly, the d_{NS}/d_S value of 0.155 was highest of all regions studied. The presence of the p4 transmembrane domain was verified in all isolates. The N-terminus

Table 2

Synonymous and non-synonymous substitution rates of different region of *Blackberry yellow vein associated virus*. Abbreviations: ORF1a: polyprotein; RdRp: RNA-dependent RNA polymerase; HSP70h: heat shock protein homolog; CP: coat protein; CPm: minor coat protein.

Region	d_{Ns}	Variance (d_{Ns})	sd (d_{Ns})	d_S	Variance (d_S)	sd (d_{Ns})	d_{Ns}/d_S
ORF1a	0.010	0.000	0.002	0.234	0.000	0.021	0.042
RdRp	0.005	0.000	0.001	0.238	0.000	0.021	0.021
P4	0.027	0.000	0.014	0.174	0.003	0.056	0.155
HSP70h	0.002	0.000	0.000	0.139	0.000	0.025	0.014
CP	0.001	0.000	0.000	0.148	0.000	0.031	0.006
CPm	0.020	0.000	0.004	0.179	0.000	0.020	0.111

d_{Ns} : number of non-synonymous substitution per non-synonymous site, averaged for all isolates; d_S : number of synonymous substitution per synonymous site, averaged for all isolates; sd: standard deviation; d_{Ns}/d_S is the ratio of non-synonymous substitution to synonymous substitution. If d_{Ns}/d_S is greater than 1, the gene is under positive selection whereas if less than 1 the gene is under negative or purifying selection.

of the HSP70 h studied showed up to 8% in nucleotide diversity, with the protein sequence diversity never exceeding 4% with a d_{Ns}/d_S of 0.014. Region 4 also covered two coding regions (C-terminus of the CP and N-terminus of CPm). The C-terminus of the CP was very conserved with limited nucleotide and amino acid diversity (6% and 3%, respectively) and d_{Ns}/d_S of 0.005. The CPm region was more diverse but still under negative selection. The maximum nucleotide and amino acid diversity was about 7% with d_{Ns}/d_S of 0.111 (Table 2).

In phylogenetic analyses, independent of the method used for inferring relationships, the isolates grouped in an erratic manner suggesting the lack of geography or host-based (wild vs. cultivated blackberry) evolution patterns (Appendix A). Absence of a dominant genotype (Appendix A) suggests that BYVaV emergence was not due to human transport of infected germplasm from a single source.

Conversely, the incongruent phylogenetic relationships suggested frequent recombination events in BYVaV populations. When region 1 was evaluated, a recombinant event was detected; a North Carolina isolate (NC6) was predicted as a recombinant between a North Carolina (NC3-like) and an Arkansas (AR8-like) isolate as major and minor donors, respectively. Similarly, one recombinant was detected in region 2 by using the same criteria: a North Carolina isolate (NC7) was predicted as a recombinant between the same North Carolina lineage as identified for region 1 (NC3-like), and a South Carolina isolate (SC3-like), as major and minor donors, respectively (Appendix A).

The possibility of recombination between the different regions studied was also evaluated after merging regions 1/2 (RNA1) and 3/4 (RNA2). There were six potential events identified, two in RNA1 and four in RNA2 (Appendix A). In all cases, there were more than six programs detecting recombination with P values always lower than 10^{-5} , adding to the validity of the results.

Virus population structures give insight into virus evolution and may facilitate understanding of disease epidemiology. In the case of closteroviruses (Moreno et al., 2008; Bertazzon et al., 2010), population structure studies provide information concerning genome size fluctuation (Tzanetakis and Martin, 2004) and evolution (Karasev, 2000; Dolja et al., 2006). Studies in the *Closteroviridae* are skewed toward members of the genera *Ampelovirus* and *Closterovirus* (Lopez et al., 1998; Kong et al., 2000; Iglesias et al., 2008; Turturo et al., 2005; Martin et al., 2009; Wang et al., 2011; Alabi et al., 2011). Unlike the above mentioned cases, population structures of criniviruses are understudied with few exceptions (Aguilar et al., 2003; Aritua et al., 2008). The most elaborate crinivirus population study to date analyzed 2277 nt/isolate or about 13% of the *Cucurbit yellow stunting disorder virus* (CYSDV) genome (Marco and Aranda, 2005).

The results of the present study show varying levels of diversity in the different regions analyzed, similar to what has been reported for *Citrus tristeza virus* (CTV; Lopez et al., 1998) and CYSDV (Marco and Aranda, 2005). In general, the nucleotide sequence is more diverse in RNA1 compared to RNA2, indicating that

criniviruses are similar to the monopartite closteroviruses where the replication-associated protein coding regions show higher nucleotide diversity compared to the coding areas found downstream (Appendix A).

Analyses of forces shaping BYVaV evolution indicate that all genome portions analyzed in this study are subjected to negative (purifying) selection, which is in agreement to what has been reported for some other closteroviruses (Alabi et al., 2011; Marco and Aranda, 2005). However, it is well documented for CTV, the better studied closterovirus, that the evolutionary forces exhibit differential selection pressures upon different proteins (Rubio et al., 2001; Martin et al., 2009). Interestingly, and comparable to SPaV (Genbank accessions EF525176 and AY948199), some BYVaV isolates show relatively high level of nucleotide diversity in one of the most conserved proteins, namely the RdRp ($d_{Ns}/d_S = 0.021$), but those changes occur mostly on the 3rd base position leading to overall amino acid preservation. Unfortunately, previous studies did not use this region in their analysis, prohibiting any comparisons between the BYVaV and other crinivirus orthologs. The selection pressure differences are also well-exemplified by different rates of selection pressure exerted on the p4-coding region and adjacent ORF coding for HSP70h, or between the partially overlapping ORFs coding for CP and CPm (Table 2). Overall, the results of our analyses indicate that regions coding for the RdRp, the N-terminus of the HSP70h protein and C-terminus of the CP, are all under stronger selection pressure than the rest of the coding regions probed in this study.

No clustering pattern was identified in phylogenetic analyses; no group-of-three-isolates were clustered together for all four regions studied (Appendix A). Such incongruence suggested that recombination among isolates is an active force in shaping BYVaV evolution. Recombination within virus populations is an important evolutionary trait in the *Closteroviridae* (Karasev, 2000) and among RNA viruses in general (Sztuba-Solinska et al., 2011). Recombination as an important evolutionary force is well supported in the case of BYVaV for two reasons, including (i) the virus is frequently found in wild or cultivated blackberries in Southeastern United States (high inoculum pressure) and (ii) the virus infects a perennial host which is exposed to several separate infection events during its lifespan. Infection of a perennial host may be an important factor for BYVaV recombination, especially when compared with CYSDV (Marco and Aranda, 2005), which infects annual hosts and did not contain any recombination signals comparable to the ones obtained for BYVaV (data not shown).

This study generates the first comprehensive information on BYVaV population structure. It will increase our ability to develop detection tests that can potentially detect all isolates of the virus, eliminating false negatives and minimizing virus escapes. Our data provide for the first time an insight into the evolution of a crinivirus that infects a perennial, woody plant. We present evidence that a stable virus population can evolve effectively through RNA recombination.

Acknowledgements

This work was funded by the US Department of Agriculture under the Specialty Crops Research Initiative (SCRI), award number 2009-51181-06022; National Clean Plant Network (NCPN), award number 10-8100-1572 and a grant from the North American Raspberry and Blackberry Association (NARBA).

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.virusres.2012.07.009>.

References

- Aguilar, J.M., Franco, M., Marco, C.F., Berdiales, B., Rodriguez-Cerezo, E., Truniger, V., Aranda, M.A., 2003. Further variability within the genus *Crinivirus*, as revealed by the determination of the complete RNA genome sequence of *Cucurbit yellow stunting disorder virus*. *Journal of General Virology* 84, 2555–2564.
- Alabi, O.J., Rwahnihi, M.A., Karthikeyan, G., Poojari, S., Fuchs, M., Rowhani, A., Rayapati, N.A., 2011. *Grapevine leafroll-associated virus 1* occurs as genetically diverse populations. *Phytopathology* 101, 1446–1456.
- Aritua, V., Barg, E., Adipala, E., Gibson, R.W., Vetten, H.J., 2008. Further evidence for limited genetic diversity among east African isolates of *Sweet potato chlorotic stunt virus*. *Journal of Phytopathology* 56, 181–189.
- Bertazzon, N., Borgo, M., Vanin, S., Angelini, E., 2010. Genetic variation and pathological properties of *Grapevine leafroll-associated virus 2* isolates. *European Journal of Plant Pathology* 127, 185–197.
- Dolja, V.V., Kreuze, F., Valkonen, J.P.T., 2006. Comparative and functional genomics of closteroviruses. *Virus Research* 117, 38–51.
- Felsenstein, J., 1985. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39, 783–791.
- Hall, T.A., 1999. BioEdit, a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symposium Series* 41, 95–98.
- Herrera-Isidrón, L., Ochoa-Sánchez, J.C., Rivera-Bustamante, R., Martínez-Soriano, J.P., 2009. Sequence diversity on four ORFs of citrus tristeza virus correlates with pathogenicity. *Virology Journal* 6, 116.
- Huang, X., Madan, A., 1999. CAP3, A DNA sequence assembly program. *Genome Research* 9, 868–877.
- Iglesias, N.G., Gago-Zachert, S.P., Robledo, G., Costa, N., Plata, M.I., Vera, O., Grau, O., Semorile, L.C., 2008. Population structure of *Citrus tristeza virus* from field Argentinean isolates. *Virus Genes* 36, 199–207.
- Karasev, A.V., 2000. Genetic diversity and evolution of closteroviruses. *Annual Review of Phytopathology* 38, 293–324.
- Kimura, M., 1980. A simple method for estimating evolutionary rate of base substitutions through comparative studies of nucleotide sequences. *Journal of Molecular Evolution* 16, 111–120.
- Kong, P., Rubio, L., Polek, M., Falk, B.W., 2000. Population structure and genetic diversity within California *Citrus tristeza virus* (CTV) isolates. *Virus Genes* 21, 139–145.
- Korber, B., 2000. HIV signature and sequence variation analysis. In: Allen, G., Rodrigo, Gerald, H., Learn (Eds.), *Computational Analysis of HIV Molecular Sequences*. Kluwer Academic Publishers, Dordrecht, Netherlands, pp. 55–72 (Chapter 4).
- Laney, A.G., Keller, K.E., Martin, R.R., Tzanetakis, I.E., 2011. A discovery 70 years in the making, characterization of Rose rosette virus. *Journal of General Virology* 92, 1727–1732.
- Lopez, C., Ayllon, M., Navas-Castillo, J., Guerri, J., Moreno, P., Flores, R., 1998. Molecular variability of 5' and 3' terminal region of *Citrus tristeza virus* RNA. *Phytopathology* 88, 685–691.
- Marco, C.F., Aranda, M.A., 2005. Genetic diversity of a natural population of *Cucurbit yellow stunting disorder virus*. *Journal of General Virology* 86, 815–822.
- Martin, R.R., Tzanetakis, I.E., Gergerich, R., Fernandez, G., Pesic, Z., 2004. *Blackberry yellow vein associated virus*, a new crinivirus found in blackberry. *Acta Horticulturae* 656, 137–142.
- Martin, S., Sambade, A., Rubio, L., Vives, M.C., Moya, P., Guerri, J., Elena, S.F., Moreno, P., 2009. Contribution of recombination and selection to molecular evolution of *Citrus tristeza virus*. *Journal of General Virology* 90, 1527–1538.
- Martin, D.P., Lemey, P., Lott, M., Moulton, V., Posada, D., Lefevre, P., 2010. RDP3, a flexible and fast computer program for analyzing recombination. *Bioinformatics* 26, 2462–2463.
- Moreno, P., Ambros, S., Abiach-Marti, M., Guerri, J., Pena, L., 2008. Citrus tristeza virus, a pathogen that changed the course of the citrus industry. *Molecular Plant Pathology* 9, 251–268.
- Martin, R.R., MacFarlane, S., Sabanadzovic, S., Quito-Avila, D.F., Poudel, B., Tzanetakis, I.E., 2012. Viruses and virus diseases of Rubus. *Plant Disease*, <http://dx.doi.org/10.1094/PDIS-04-12-0362-FE>.
- Nei, M., Gojobori, T., 1986. Simple methods for estimating the numbers of synonymous and non-synonymous nucleotide substitutions. *Molecular Biology and Evolution* 3, 418–426.
- Poudel, B., 2011. Epidemiological studies on *Blackberry yellow vein associated virus* and *Blackberry chlorotic ringspot virus*. MS Thesis, University of Arkansas, Fayetteville, AR, 88 pp.
- Roy, A., Manjunath, K.L., Bransky, R.H., 2005. Assessment of sequence diversity in the 5'-terminal region of *Citrus tristeza virus* from India. *Virus Research* 113, 132–142.
- Rubio, L., Ayllon, M.A., Kong, P., Fernandez, A., Polek, M., Guerri, J., Moreno, P., Falk, B., 2001. Genetic variation of *Citrus tristeza virus* isolates from California and Spain, evidence of mixed infections and recombination. *Journal of Virology* 75, 8054–8062.
- Susaimuthu, J., Tzanetakis, I.E., Geregrich, R.C., Martin, R.R., 2006. Yellow vein-affected blackberries and presence of a novel *crinivirus*. *Plant Pathology* 55, 607–613.
- Susaimuthu, J., Geregrich, R.C., Bray, M.M., Clay, K.A., Clark, J.R., Tzanetakis, I.E., Martin, R.R., 2007. Incidence and ecology of *Blackberry yellow vein associated virus*. *Plant Disease* 91, 809–813.
- Susaimuthu, J., Tzanetakis, I.E., Gergerich, R.C., Kim, K.S., Martin, R.R., 2008. Viral interactions lead to decline of blackberry plants. *Plant Disease* 92, 1288–1292.
- Sztuba-Solinska, J., Urbanowicz, A., Figlerowicz, M., Bujarski, J.J., 2011. RNA-RNA recombination in plant virus evolution and replication. *Annual Review of Phytopathology* 49, 415–443.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S., 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28, 2731–2739.
- Turturo, C., Saldarelli, P., Yafeng, D., Digiario, M., Minafara, A., Savino, V., Martelli, G.P., 2005. Genetic variability and population structure of *Grapevine leafroll-associated virus 3* isolates. *Journal of General Virology* 86, 217–224.
- Tzanetakis, I.E., Martin, R.R., 2004. Complete nucleotide sequence of a strawberry isolate of *Beet pseudo-yellows virus*. *Virus Genes* 28, 239–246.
- Tzanetakis, I.E., Postman, J.D., Martin, R.R., 2005. A member of the *Closteroviridae* from mint with similarities to all three genera of the family. *Plant Disease* 89, 654–658.
- Tzanetakis, I.E., Susaimuthu, J., Gergerich, R., Martin, R.R., 2006. Nucleotide sequence of *Blackberry yellow vein associated virus*, a novel member of *Closteroviridae*. *Virus Research* 116, 196–200.
- Tzanetakis, I.E., Halgren, A., Mosier, N., Martin, R., 2007. Identification and characterization of *Raspberry mottle virus*, a novel member of the *Closteroviridae*. *Virus Research* 127, 26–33.
- Wang, J., Sharma, A.M., Duffy, S., Almeida, R.P., 2011. Genetic diversity of 3' terminal 4.7-kb region of *Grapevine leafroll-associated virus 3*. *Phytopathology* 101, 445–450.
- Zuckerland, E., Pauling, L., 1965. Evolutionary divergence and convergence in proteins. In: Bryson, V., Vogel, H.J. (Eds.), *Evolving Genes and Proteins*. Academic Press, New York, pp. 97–166.